References:

1. William P. J and Douglas K. A., (2008), Biologically Active Natural Products of the Genus *Callicarpa*, Current Bioact. Compound. 4(1),15-32.
2. Qadrie Z, Rajkapoor B. and Kavimani S. (2013) Acute and Subacute Toxicity of *Callicarpa linata* leaf Extracts in Rats. International Journal of Pharmaceutical & Research Sciences. 2 (11), 1095-1104.
3. Amin M., Ibrahim S. and Santoni A. (2015) Preliminary phytochemical and HPLC screening of triterpenoids fraction from bark extract of *Callicarpa Arborea* Roxb. Journal of Chemical and Pharmaceutical Research. 7(10),848-851.
4. Cantrell C. L., Klun J. A., Bryson C. T., Kobaisy M. and Duke S. O. (2005) Isolation and Identification of Mosquito Bite Deterrent Terpenoids from Leaves of American (*Callicarpa americana*) and Japanese (*Callicarpa japonica*) Beautyberry. J. Agric. Food Chem. 53, 5948-5953.
5. William P. J., Tatiana L. E., Qiuwen M. Hyee B. C., Djaja D. S., Geoffrey A. C., Steven M. S., and Douglas A. K. (2007) Cytotoxic Constituents from the Fruiting Branches of Callicarpa americana Collected in Southern Florida. J. Nat. Prod. 70, 372-377.
6. Tu Y., Sun L., Guo M., and Chen W. (2013). The medicinal uses of Callicarpa L. in traditional Chinese medicine: an ethnopharmacological, phytochemical and pharmacological review. J Ethnopharmacol. 3;147(3),680-682.
7. Yung A. K., S H. S., Jong H. P., Seung H. K. Lee Y. K. and Young C.K. (2005). In Vitro Neuroprotective Activities of Phenyl ethanoid Glycosides from Callicarpa dichotoma. Letter- Planta Med. 71, 778 – 780.
8. Verma V. K., Siddiqui N.S. and Aslam M. (2012). A new Kaurane Diterpene from the leaves of *Callicarpa Macrophylla* Vahl . International Research Journal of Pharmacy. 3 (5), 178-179.
9. Qadrie Z., Rajkapoor B. and Kavimani S. (2013). Antioxidant Activity of Ethanolic Extracts of *Callicarpa Linata* Leaf. *International Journal of Pharmacological Research.* 5(6), 129-131.

10. Chung P.Y. , Chung L. Y. and Navaratnam P. (2014) . Potential targets by entacyclic triterpenoids from *Callicarpa farinosa* against methicillin-resistant and sensitive Staphylococcus aureus. Fitoterapia. 94, 48–54.
11. Lallianchhunga M.C., Devi L. I., Lalmuanthanga C., Lalchhandama C., Subudhi P.K. and Ali A. M. [2016] *in vitro* antioxidant activity of methanolic extract of *callicarpa arborea* leaves. World Journal of Pharmaceutical Research.5(4), 2097-2102.
12. Munir A. A. (1982). A Taxonomic Revision of the Genus Callicarpa L. (Verbenaceae) in Australia. J. Adelaide Bot. Gard. 6(1), 5-39.
13. Harshitha K ., Nishteswar K. and Harisha C.R.(2013). Pharmacognostical and Preliminary Phytochemical Investigations on Different Parts of Bulbophyllum Neilgherrense Wight. -An Orchid Used in Folk Medicine. Global Journal of Research on Medicinal Plant and Indigenous Medicines. 2(4), 259-269.
14. Devi Priya M. and E.A. Siril (2013). Pharmacognostic Studies on Indian Madder (Rubia cordifolia L.). Journal of Pharmacognosy and Phytochemistry. 1(5)112-119.
15. Suresh kumar C.A., Varadharajan R., Muthumani P., Meera R., Devi P. and Kameswari B. (2009). Pharmacognostic and Preliminary Phytochemical Investigations on the stem of *Saccharum spontaneum.* J. Pharm. Sci. & Res. 1(3), 129-136.
16. Rajendran R., Hemalatha S., Akasakalai K., MadhuKrishna C.H., Sohil B., Vittal and Meenakshi R. S. (2009). Hepatoprotective activity of Mimosa pudica leaves against Carbontetrachloride induced toxicity. Journal of Natural Products Vol. 2(2009):116-122.
17. Roy C. K. and Das A. K. (2010) Effect of *Psidium Guajava* Linn. Methanolic Leaf Extract On Hepatoprotection. Journal of Pharmaceutical And Biomedical Sciences. 1 (03),01-04.
18. Panchal C.V., Sawale J. A., Poul B. N. and Khandelwal K.R. (2013). Hepatoprotective activity of lagenaria siceraria (molina) standley fruits against paracetamol induced hapatotoxicity in mice. IJPSR, Vol. 4(1): 371-377.

19. Hurkadale P. J., Shelar P.A. , Palled S.G. , Mandavkar Y.D. and Khedkar A. S. (2012). Hepatoprotective activity of Amorphophallus paeoniifoliustubers against paracetamol-induced liver damage in rats. Asian Pacific Journal of Tropical Biomedicine. S, 238-242.

20. Nwala C. O., Akaninwor J. O. and Monanu M. O. (2013). Phytochemical Screening and wound healing activities of Extracts of Jatropha Curcas leaf formulated in a Simple ointment Base. International Journal of Engineering Science Invention. 2 (6), 72-75.
21. Shafiuddin M., Khan A. And Ali S. (2009).Wound Healing Activity of Traditional Herbal Formulation. Int. J. Chem. Sci. 7(2), 639-643.
22. Vijendren S., Jothi S., Rajandran K. and Muruganandham V. (2012). Assessment of In-vitro Wound Healing Activity of the *tinospora crispa* Extracts. International Journal of Pharmacology and Pharmaceutical Technology. 1(05), 01-05.
23. AbdullaM. A., Ahmed K.A. and Abu-Luhoom F.M.(2010), Role of Ficus deltoidea extract in the enhancement of wound healing in experimental rats. Biomedical Research. 21 (3), 241-245.
24. Govindam S., Kuchi M., Balekari U. and Rani G. S. (2011). Screening of Wound Healing Effect of Bark of *Barringtonia Asiatica* . International Journal of Pharmacology Research. 1(1),26-31.
25. Sarkar M., Das G., Pathak S.K., Maitra S. And Samanta S. (2013). Evaluation of In Vivo Wound Healing and In Vitro Antibacterial Activities of The Different Extract of Leucas Indica Linn. International Journal of Pharmacy and Pharmaceutical Sciences. 5(3),333-340.
26. Ekpo M., Mbagwu H., Jackson C. and Eno M. (2011). Antimicrobial and Wound Healing Activities of Centrosema Pubescens (Leguminosae). Journal of *Pharmacy* and Clinical Sciences. 1.,01-06.
27. Nwala C. O., Akaninwor J. O. and Monanu M. O. (2013) Use of Extracts of Jatropha Curcas Leaf Formulated in A Simple Ointment Base in Wound Healing Activities: How Safe Is It? International Journal of Engineering Science Invention. 2(6),53-57.

28. Morshed A.J.M., (2013). A Survey of Medicinal Plants as Regards to their Uses by the Tribal Practitioners in the Chittagong Hill Tracts of Bangladesh to Check Various Diseases. Hamdard Medicus. 56(3).
29. Devi Prasad A.G. and Shyma T.B.(2013) .Medicinal plants used by the tribes of Vythiri taluk, Wayanad district (Kerala state) for the treatment of human and domestic animal ailments, JMPR 7(20):1439-1451.
30. Ratheesh Narayanan M.K., Mithunlal S., Sujanapal P., Anil Kumar N., Sivadasan M. and Alfarhan A.H. (2011). Ethnobotanically important trees and their uses by Kattunaikka tribe in Wayanad Wildlife Sanctuary, Kerala, India, JMPR 5(4):604-612.
31. Jothy S. L., Yeng C. and Sasidharan S. (2013). Chromatographic and Spectral Fingerprinting of *Polyalthia longifolia*, a Source of Phytochemicals BioResources 8(4), 5102-5119.
32. Arnóbio A., Silva-J., Silvana M.Z. and Matheus de Freitas F.P. (2017). Medicinal Plants for the Treatment of Local Tissue Damage Induced by Snake Venoms: An Overview from Traditional use to Pharmacological Evidence. Evidence-Based Complementary and Alternative Medicine.01-52.